

Taxonomy of *Juniperus deppeana* varieties and formas based on nrDNA (ITS), petN-psbM, trnS-trnG, trnD-trnT, trnL-trnF sequences

Robert P. Adams

Biology Department, Baylor University, Box 97388, Waco, TX 76798, USA
Robert_Adams@baylor.edu

Andrea E. Schwarzbach

Department of Biomedicine, University of Texas at Brownsville, Brownsville, TX 78520, USA.

ABSTRACT

Juniperus deppeana has numerous disjunct populations that include four taxonomic varieties and three forms. All four varieties and two forms of *Juniperus deppeana* from the southwest United States, Mexico and Guatemala were analyzed by sequencing nrDNA (ITS), petN-psbM, trnS-trnG, trnD-trnT, trnL-trnF. A Bayesian tree gave support for clades of var. *deppeana*, NM, var. *patoniana*, and var. *gamboana*. However, several clades with high support contain mixtures of different varieties and forms. A minimum spanning network based on 91 mutational events (MEs) showed that the varieties and forms are extremely closely related, differing by only 1 to 2 bp (out of 4411 bp). The taxon with the largest differentiation was var. *deppeana*, Sacramento Mtns., NM that differed by 4 MEs from the Oak Creek canyon, AZ individuals. The lack of variation among *J. deppeana* taxa may be due to the mixing of populations during the Wisconsin glacial maximum (70,000 - 13,000 ybp) when life zones descended about 800 m. Published on-line: www.phytologia.org *Phytologia* 95(2): 161-166 (May 1, 2013).

KEY WORDS: *Juniperus deppeana* varieties, Cupressaceae, DNA, nrDNA (ITS), petN-psbM, trnS-trnG, trnD-trnT, trnL-trnF, systematics, geographic variation, taxonomy.

Juniperus deppeana Steudel has trunk bark that exfoliates in quadrangular plates, thus the common name 'alligator bark' juniper. *Juniperus deppeana* is part of the serrate leaf margined species of the western hemisphere (Adams, 2011) and is widely distributed in the southwestern US, Mexico and northern Guatemala (Fig. 1). Putative *Juniperus d. f. sperryi*, once known only from the type locality in the Davis Mtns., TX, has been found in Arizona and New Mexico (Fig. 1). However, the bark characters seem to be controlled by only a few genes thus furrowed bark may have arisen independently in western Mexico and the southwestern United States (e.g., *J. d. var. patoniana* and *J. d. f. sperryi*). Whether the furrowed bark trees of Mexico are related to *J. d. f. sperryi* is not well understood.

The first systematic treatment of the serrate leaf-margined junipers was by Martinez (1963) who recognized *J. deppeana* Steudel. var. *deppeana* (checked bark, (3)4-5(6) seeds/cone,




Figure 1. Distribution of *J. deppeana*.

J. d. var. pachyphlaea (Torrey) Mart. (checkered bark, (1)2-4(5) seeds/cone), *J. d. var. robusta* Mart. (checkered bark, (1)2-3(-6) seeds/cone), *J. d. var. zacatecensis* Mart. (checkered bark, 1-4(-7) seeds/cone), *J. patoniana* Mart. (laced bark, (1)2-3(-6) seeds/cone, and *J. gamboana* Mart. (checkered bark, 1(2) seeds/cone) (Fig. 2).




Figure 2. Variation in bark exfoliation among *J. deppeana* varieties and forms.

Zanoni and Adams (1976, 1979) and Adams, Zanoni and Hogge (1984), using morphology and essential oils, generally agreed with Martinez's treatment, except *J. patoniana* was reduced to *J. d. var. patoniana* (Mart.) Zanoni. Additional studies (Adams and Nguyen, 2005; Adams et al. 2007) have further clarified geographical variation in *J. deppeana*.

Recently, Adams and Schwarzbach (2006) recognized *J. gamboana* as *J. deppeana* var. *gamboana* (Mart.) R. P. Adams and *J. deppeana* var. *zacatecensis* as *J. deppeana* f. *zacatecensis* (Mart.) R. P. Adams. Adams and Schwarzbach (2011) found *J. deppeana* and var. *gamboana* to be a clade, sister to *J. ashei*, *J. saltillensis* and *J. zanonii* (Fig. 3). However, the other *J. deppeana* varieties and forms were not included in their study.

The focus of the present study was to examine relationships among all the recognized (Adams 2011) varieties and forms of *J. deppeana* (except the very minor variant f. *elongata*, Adams 2011) using data obtained from sequencing of nrDNA (ITS), petN-psbM, trnS-trnG, trnD-trnT and trnL-trnF.

MATERIALS AND METHODS

Specimens used in this study: *J. deppeana* var. *deppeana*, Adams 10539-10541, El Chico National Park, Hidalgo, MX; Adams 7632-7634, Sacramento Mtns., e of Alamogordo, NM, USA; Adams 10640-10642, Oak Creek Canyon-Flagstaff, AZ; *J. deppeana* var. *gamboana*, Adams 6863-6867, Comitán, Chiapas, MX; *J. deppeana* var. *patoniana*, Adams 6836-6839, km 152, w. of Durango (city), Durango, MX (P); *J. deppeana* var. *robusta*, Adams 10255-10256, w of La Ciudad, Durango, MX; *J. deppeana* f. *sperryi*, Adams 10626, Bridge Spring, Davis Mtns., TX, USA; Adams 11312, Munds Mtn., AZ; *J. deppeana* f. *zacatecensis*, Adams 6840-6842, 18 km w. Sombrette, Zacatecas, MX; *J. virginiana*, Adams 10231-10232, Knoxville, TN, USA. Voucher specimens are deposited at BAYLU herbarium, Baylor University.




Figure 3. Bayesian tree of the serrate *Juniperus* of North America. Numbers at the branch points are posterior probabilities (as percent).

One gram (fresh weight) of the foliage was placed in 20 g of activated silica gel and transported to the lab, thence stored at -20° C until the DNA was extracted. DNA was extracted from juniper leaves by use of a Qiagen mini-plant kit (Qiagen, Valencia, CA) as per manufacturer's instructions.

Amplifications were performed in 30 μ l reactions using 6 ng of genomic DNA, 1.5 units Epi-Centre Fail-Safe Taq polymerase, 15 μ l 2x buffer E (petN, trnD-T, trnL-F, trnS-G) or K (nrDNA) (final concentration: 50 mM KCl, 50 mM Tris-HCl (pH 8.3), 200 μ M each dNTP, plus Epi-Centre proprietary enhancers with 1.5 - 3.5 mM $MgCl_2$ according to the buffer used) 1.8 μ M each primer. See Adams, Bartel and Price (2009) for the ITS and petN-psbM primers utilized. The primers for trnD-trnT, trnL-trnF and trnS-trnG regions have been previously reported (Adams and Kauffmann, 2010).

The PCR reaction was subjected to purification by agarose gel electrophoresis. In each case, the band was excised and purified using a Qiagen QIAquick gel extraction kit (Qiagen, Valencia, CA). The gel purified DNA band with the appropriate sequencing primer was sent to McLab Inc. (San Francisco) for sequencing. Sequences for both strands were edited and a consensus sequence was produced using Chromas, version 2.31 (Technelysium Pty Ltd.) or Sequencher v. 5 (genecodes.com). Sequence datasets were analyzed using Geneious v. 5.4 (Drummond et al. 2011), the MAFFT alignment program and the PAUP* program, version 4.0b10 (Swofford 2003) for neighbor joining, parsimony, and maximum likelihood tree searches. Further analyses utilized the Bayesian analysis software Mr. Bayes v.3.1

sufficient, by themselves, to warrant the recognition of a new variety from NM. Additional research on geographic variation in leaf terpenes and morphology is in progress so as to examine differentiation in the southwestern US.

CONCLUSION

The mixtures of various taxa within clades may be due to ancient climate and past distributions of *J. deppeana*. Wells (1966), using data from rat middens from the Big Bend of Trans-Pecos, Texas, concluded that during the Wisconsin (70,000 - 13,000 ybp) life zones descended about 800 m leading to the formation of a pinyon-juniper woodland in the present Chihuahuan desert between the Big Bend of Trans-Pecos, Texas and the city of Del Rio. Assuming that the effects of glaciation were mediated southward into Mexico so that life zones descended only a few hundred meters in Hidalgo, it appears that most of the now disjunct populations of *J. deppeana* may have once been connected in a nearly continuous population of distribution around the Chihuahuan desert (Fig. 5). It is likely that desert peaks within the area concerned also supported stands of *J. deppeana*. Wisconsin populations would have become spatially separated as dryer, warmer climate developed during the Holocene (past 13,000 y). Of course, the Wisconsin was only the most recent of several pluvial events during the Pleistocene, spanning 1.8 my (Flint, 1971). It is likely that during any one (or several) of these pluvial events, *Juniperus deppeana* occupied lower elevation and more southward habitats, leading to more contiguous populations in Mexico and the southwestern United States. If divergent populations (or varieties) became sympatric during the Wisconsin, this would have facilitated infra-specific crossing. This may account for the large genetic variation within some populations. In addition, the millennia of continuous populations could explain the lack of differentiation between the recently (Holocene) geographically isolated populations.




Figure 5. Possible range of *J. deppeana* during the Wisconsin glacial maximum (based on Wells, 1966). The present day disjunct populations were likely continuous in the foothills around the Chihuahuan desert during the Wisconsin.

ACKNOWLEDGEMENTS

Thanks to Tom Zanoni (NYBG) for providing slides of tree barks and assistance in field work, and Tonnie Yanke for lab assistance. This research was supported in part with funds from NSF grant DEB-316686 (A. Schwarzbach and R. P. Adams) and funds from Baylor University.

LITERATURE CITED

- Adams, R. P. 1975. Statistical character weighting and similarity stability. *Brittonia* 27: 305-316.
- Adams, R. P. 2011. The junipers of the world: The genus *Juniperus*. 3rd ed. Trafford Publ., Victoria, BC.
- Adams, R. P., J. A. Bartel and R. A. Price. 2009. A new genus, *Hesperocyparis*, for the cypresses of the new world. *Phytologia* 91: 160-185.
- Adams, R. P. and S. Nguyen. 2005. Infra-specific variation in *Juniperus deppeana* and f. *sperryi* in the Davis Mountains, Texas: Variation in leaf essential oils and Random Amplified Polymorphic DNAs (RAPDs). *Phytologia* 87:96-108.
- Adams, R. P. and A. E. Schwarzbach. 2006. Intraspecific adjustments in *Juniperus deppeana* (Cupressaceae) *Phytologia* 88: 227-232.
- Adams, R. P. and A. E. Schwarzbach. 2011. DNA barcoding a juniper: the case of the south Texas Duval county juniper and serrate junipers of North America. *Phytologia* : 146-154.
- Adams, R. P. and A. E. Schwarzbach. 2012. Taxonomy of the multi-seeded, entire leaf taxa of *Juniperus* section *Sabina*: Sequence analysis of nrDNA and four cpDNA regions. *Phytologia* 94: 350-368.
- Adams, R. P., T. A. Zanoni and L. Hogge. 1984 Analyses of the volatile oils of *Juniperus deppeana* and its infraspecific taxa: chemosystematic implications. *Biochem. Syst. and Ecol.* 12:23-28.
- Adams, R. P., A. E. Schwarzbach, S. Nguyen, and J. A. Morris. 2007. Geographic variation in *Juniperus deppeana*. *Phytologia* 89: 132-150.
- Axelrod, D. I. 1958. Evolution of the Madro-Tertiary geoflora. *Bot.Rev.* 24: 433- 509.
- Drummond A. J., B. Ashton, S. Buxton, M. Cheung, A. Cooper, C. Duran, M. Field, J. Heled , M. Kearse, S. Markowitz, R. Moir, S. Stones-Havas, S. Sturrock, T. Thierer and A. Wilson. 2011. Geneious v 5.4, Available from <http://www.geneious.com/>
- Flint, R. F. 1971. Glacial and quaternary geology. John Wiley and Sons, NY.
- Martinez, M. 1963. Las pinaceas mexicanas. Tercera edicion. Universidad Nacional Autonoma de Mexico. Mexico City.
- Posada, D. and K. A. Crandall. 1998. MODELTEST: testing the model of DNA substitution. *Bioinformatics* 14: 817-818.
- Ronquist, F. and J. P. Huelsenbeck. 2003. MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19: 1572-1574.
- Swofford, D. L. 2003. PAUP*: phylogenetic analysis using parsimony (*and other methods), version 4. Sinauer Associates, Sunderland, MA.
- Veldman, D. J. 1967. Fortran programming for the behavioral sciences. Holt, Rinehart and Winston Publ., NY.
- Wells, P. V. 1966. Late Pleistocene vegetation and degree of pluvial climatic change in the Chihuahuan desert. *Science* 153: 970-975.
- Zanoni, T. A. and R. P. Adams. 1976. The genus *Juniperus* (Cupressaceae) in Mexico and Guatemala: Numerical and chemosystematic analysis. *Biochem. Syst. Ecol.* 4: 147-158.
- Zanoni, T. A. and R. P. Adams. 1979. The genus *Juniperus* (Cupressaceae) in Mexico and Guatemala: Synonymy, Key, and distributions of the taxa. *Bol. Soc. Bot. Mexico* 39: 83-121.